

The Peculiar 2004 Superoutburst in the Helium Dwarf Nova, 2003aw

Daisaku NOGAMI¹, Berto MONARD², Alon RETTER³, Alex LIU⁴, Makoto UEMURA⁵, Ryoko ISHIOKA⁶,
Akira IMADA⁶, and Taichi KATO⁶

¹*Hida Observatory, Kyoto University, Kamitakara, Gifu 506-1314, Japan (nogami@kwasan.kyoto-u.ac.jp)*

²*Bronberg Observatory, PO Box 11426, Tiegerpoort 0056, South Africa*

³*Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Lab, University Park, PA, 16802-6305, USA*

⁴*Norcape Observatory, PO Box 300, Exmouth 6707, Australia*

⁵*Departamento de Física, Facultad de Ciencias Físicas y Matemáticas, Universidad de Concepción, Casilla 160-C, Concepción, Chile*

⁶*Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan*

(Received 2004 0; accepted 2004 0)

Abstract

We conducted a time-resolved photometric campaign of the helium dwarf nova, 2003aw in 2004 May–June. 2003aw stayed at 14.7–15.7 mag for the first several days during this campaign, which is the plateau phase of this superoutburst. This variable then rapidly decayed to ~ 18.0 mag, still brighter by about 2 mag than its quiescence magnitude, and maintained this brightness for about 20 days, having short flares of ~ 2 mag. A long fading tail followed it. We detected superhumps with a period of $0.02357(4)$ d [= 2036(3) s] during the plateau phase. The whole light curve of the superoutburst in 2003aw, taking into account the present data and those in the literature, perfectly resembles that of the 1996–1997 superoutburst of the peculiar WZ Sge-type hydrogen-rich dwarf nova, EG Cnc.

Key words: accretion, accretion disks — stars: dwarf novae — stars: individual (SN 2003aw) — stars: novae, cataclysmic variables — stars: oscillations

1. Introduction

Interacting binaries consisting of two white dwarfs (IBWDs) are called AM CVn stars (Solheim 1995; Warner 1995a). Helium gas transferred from the less massive white dwarf forms an accretion disk around the primary. There are 11 confirmed members including 2003aw in this class, 10 of which (and 2 candidates) are listed in table 1 in Woudt, Warner (2003), and SDSS J1240–01 is the most recently discovered (Roelofs et al. 2004). They have three activity groups, mainly depending on the mass-transfer rate $\dot{M}$, like hydrogen-rich cataclysmic variables (CVs; for a review, Warner 1995b). They are, from the high- $\dot{M}$ systems, 1) stable systems with a hot accretion disk, corresponding to nova-like in CVs, 2) systems having outbursts, corresponding to dwarf novae, 3) stable systems with a cold disk. 2003aw is the 6th member of the helium dwarf novae.

This star was discovered as a supernova (therefore named 2003aw) by Wood-Vasey et al. (2003) from the unfiltered NEAT image taken with the Palomar 1.2m Schmidt telescope. The magnitude at the discovery on 2003 February 6 was 17.8, and this ‘supernova’ was seen also in the images taken on 2003 February 10 and 19. Chornock, Fillipenko (2003) obtained an optical spectrum, which had a quite blue continuum with weak He I emission lines and Ca II H and K absorption at nearly zero redshift. They commented that 2003aw is not a supernova, but rather is a hydrogen-deficient dwarf nova

similar to that proposed for 1998di = KL Dra (see Jha et al. 1998).

Woudt, Warner (2003) carried out intensive high-speed photometry of 2003aw in an active phase between 2003 February 28 and June 9. They divided their observations into three states. In the high state, 2003aw was in $V = 17.6 - 19.0$, and gave rise to a flare reaching $V = 16.5$. In the intermediate state and quiescence, 2003aw was at about $V = 19.6$ and at around $V = 20.3$, respectively. Several curious features were revealed by them, such as superhumps with a period of $2041.5 (\pm 0.3)$ s, brightness cycles over 0.4 mag with a period of ~ 16 h during the high and intermediate state, and sidebands to the principal frequencies having a constant frequency difference from the superhump harmonics.

Being informed by P. Woudt on 2004 May 17 that 2003aw was in outburst at $V \sim 15$ mag, which is at least 1.4 mag brighter than the maximum during the 2003 February/March outburst [vsnet-alert 8131, see Kato et al. (2004c) on VSNET], a photometric observation campaign was immediately started by the VSNET collaboration team. We report the results in this paper.

2. Observation

The observations were carried out at four sites, Tiegerpoort in South Africa (BM), Exmouth in Australia (AL), Concepcion in Chile (MU), and Kyoto in Japan (RI, AM, and TK). Table 1 summarizes the log of the obser-

Table 1. Log of observations.

Date	HJD-2453000 Start–End	Exposure Time (s)	Useful Frames	Mean Mag.	Comparison Star*	Site [†]	Sky Condition
2004 May	18 143.960–144.058	60	86	15.7(0.1)	1	A	thin clouds
	144.005–144.019	30	35	15.2(0.8)	1	B	
	144.173–144.329	28	440	15.2(0.1)	1	C	
	19 145.179–145.304	28	353	15.3(0.1)	1	C	thin clouds
	20 145.953–146.060	60	93	14.7(0.1)	1	A	
	146.182–146.287	28	202	15.7(0.1)	1	C	
	22 147.956–148.057	60	90	16.6(0.2)	1	A	
	24 149.974–149.979	30	4	16.1(0.7)	1	B	
	25 150.960–151.057	60	61	17.6(0.3)	1	A	
	26 152.169–152.260	28	254	16.4(0.2)	1	C	
	27 152.975–153.051	90	50	18.3(0.3)	2	A	
	28 153.959–154.068	90	72	18.1(0.3)	2	A	
	29 154.953–155.057	90	66	18.1(0.3)	2	A	
June	5 162.447–162.454	30	8	18.6(0.9)	1	D	thin clouds
	6 163.457–163.463	30	9	18.1(0.6)	1	D	
	9 166.181–166.203	28	59	16.1(0.1)	1	C	
	166.432–166.536	30	81	17.0(0.6)	1	D	
	12 169.445–169.506	30	24	17.5(0.7)	1	D	
	29 186.442–186.447	30	3	18.0(0.9)	1	D	

*1: 2UCAC 29829807, 11.3 mag, 2: 2UCAC 29829819, 16.1 mag

[†]A: 30cm Tel. + SBIG ST-7E (Exmouth, Australia), B: 30cm Tel. + SBIG ST-7E (Kyoto, Japan), C: 32cm Tel. + SBIG ST-7E (Tiegerpoort, South Africa), D: 30cm Tel. + Pictor 416XT (Concepcion, Chile)

vations and the instruments.

The differential magnitudes of 2003aw were measured, assuming the unfiltered CCD magnitudes of the nearby comparison stars, 2UCAC 29829807 and 2UCAC 29829819, to be 11.3 and 16.1, respectively, based on the UCAC2 magnitude (see Zacharias et al. 2004 regarding UCAC2). The UCAC bandpass is 579–642 nm, between V and R_c . The error in the comparison magnitude is expected to be within 0.5 mag.

Heliocentric corrections to the observation times were applied before the following analysis.

3. Results

The long-term light curve is drawn in figure 1. At the first stage of our campaign, 2003aw was recorded to be at 15.1–15.7 mag with a brightening to 14.7 mag on May 20. Even considering the zero-point uncertainty of the magnitude and the color, this maximum magnitude far exceeds the brightest magnitude at a brightening ($V = 16.5$) in the ‘high’ state during the 2003 February/March outburst (Woudt, Warner 2003). The detection of the superhumps (described later) indicates that this phase corresponds to the plateau phase of the superoutburst in AM CVn stars and SU UMa-type dwarf novae. It is unclear how long the plateau phase lasted before our observations started.

After the decline from the plateau phase, 2003aw kept its brightness around 18.0 mag, besides occasional flares reaching ~ 15.8 mag, for ~ 20 days. During this phase, 2003aw was still brighter by about 2.0 mag than in qui-




Fig. 1. Long-term light curve of the 2004 May outburst of 2003aw. The abscissa is HJD – 2453000, and the ordinate is CCD magnitude. The open circles are the CCD observations reported to VSNET. The error of these observations depends on the target brightness and the sky condition, but is smaller than 0.4 mag. The filled squares are average magnitudes of each run of our data. A quite bright state was recorded for several days at the start of our campaign. Then 2003aw stayed around 18 mag for about 20 days, occasionally having sudden, short flares. After this state, 2003aw gradually went down to quiescence.




Fig. 2. Short-term light curves of the long runs during the plateau phase. Stable superhumps with a small amplitude are seen in panels (b) and (c). The superhump phase of the abscissa is adjusted using the data shown in these panels. There are large amplitude (~ 0.5 mag) modulations in panel (a), which are possibly related to the superhumps. There is no clear evidence of the superhump in panels (d) and (e).

escence, which corresponds to the high state in Woudt, Warner (2003). Judging from our observations and those reported to VSNET (see figure 1), the duration of the flares is 1 day, or so, but a periodicity of the flare can not be supported nor rejected from the available sparse data. This behavior is completely consistent with that during the high state in the 2003 February/March active phase (Woudt, Warner 2003).

A gradual fading to quiescence [the intermediate state defined by Woudt, Warner (2003)] followed the high state. We could not, however, continue the observations well during and after this state with our 30cm telescopes.

Figure 2 shows the enlarged light curves of each long run during the plateau phase. There exist modulations with variable shapes, amplitudes, and timescales. Those shown in figures 2a have a large amplitude up to 0.5 mag and an unstable timescale. We see small, stable modulations in figures 2b and 2c, which are interpreted as superhumps. A rapid brightening with a rate of 4.4 mag d^{-1} is displayed in figure 2d, but 2003aw decayed by 0.9 mag during the following 3 hours by the start of the run of figure 2e.

We performed period analyses by the Phase Dispersion Minimization (PDM) method (Stellingwerf 1978) and the Fourier transformation for all the data in figures 2b and 2c containing the stable superhumps. Before these analyses, a pre-whitening removing variations with timescales longer than hours was processed. The resultant Θ diagram and power spectrum are exhibited in figure 3. The best estimated superhump period is $0.02357 (\pm 0.00004) \text{ d}$ [= $2036 (\pm 3) \text{ s}$]. The error of the period was estimated using the Lafler-Kinman class of methods, as applied by Fernie (1989). This period is different by $\sim 2 \sigma$ from the value [$2041.5 (\pm 0.3) \text{ s}$] derived by Woudt, Warner (2003). This difference can not be considered statistically significant, though a similar difference in the superhump period in the different outburst states were observed in another AM CVn star, V803 Cen (Kato et al. 2004b). Including the data shown in figures 2a and 2d did not improve the




Fig. 3. Theta diagram of the PDM analysis and power spectrum, using the data shown in figure 2b and 2c, after removing variations with timescales longer than hours. Panels (b) and (d) show the results in panels (a) and (c), respectively, zoomed on the frequency range around the superhump frequency. Both analyses indicate the superhump period of $0.02357(4) \text{ d}$ [= $42.44(7) \text{ cycle d}^{-1} = 2036(3) \text{ s}$].

period determination.

We could not detect firm evidence of any sidebands, which would be (at least partly) due to short coverages and low signal-to-noise ratios of our data.

The averaged superhump profiles are exhibited in figure 4. There is a hint of a dip after the superhump maximum on 2004 May 18 (figure 4a). The same phenomenon was observed by Woudt, Warner (2003) during the high state. On this day, the superhumps had a shape consisting of a rapid rise and a slow decline, which is typically seen in hydrogen-rich SU UMa-type dwarf novae. On the next day, 2004 May 19, the superhump had evolved into a symmetric shape (figure 4b).

The timings of the superhump maxima were extracted




Fig. 4. The superhump profile folded with the superhump period derived in this work and averaged. Panels (a) and (b) use the data shown in figures 2(b) and 2(c), respectively. There is a hint of a dip around phase 0.1 in (a) superposed on a typical superhump shape with a rapid rise and a slow decay. During the following ~ 40 superhump cycles, the superhump shape became rather symmetric.

by fitting the average superhump light curve in figure 4, and we tried to check the change of the superhump period between HJD 2454144.173 and 2453145.304. The superhump period, however, stayed constant during this period.

A periodicity attributable to the orbital modulation could not be detected from any of our runs.

4. Discussion

Our observations of 2003aw revealed the existence of the plateau phase of the superoutburst, which is brighter by ~ 5.0 mag than its quiescent magnitude, and the presence of superhumps with a period of 2036(3) s during this phase. Following this phase, the star sharply decayed with a rate over 1 mag d^{-1} and stayed around 18.0 mag in the high state for ~ 20 days, having short flares with amplitudes of ~ 2.0 mag. This variable gradually faded after that. Woudt, Warner (2003) are considered to have missed the plateau phase, and have steadily observed the high state and the fading tail.

The whole outburst light curve of 2003aw, taking into account the observations by us and Woudt, Warner (2003), has strong resemblance to the superoutburst of the peculiar WZ Sge-type dwarf nova, EG Cnc, in 1996–1997 (Patterson et al. 1998; Kato et al. 2004a), in terms of the first quite bright state (the plateau phase in 2003aw, and the main superoutburst in EG Cnc), the following repetitive-outburst state still brighter than the quiescence (the high state in 2003aw, and the rebrightening phase in EG Cnc), and the long fading tail to the quiescence (the intermediate state in 2003aw, and the long fading tail in EG Cnc). Superhumps were observed throughout these states in both stars.

The thermal-tidal disk-instability models for the AM CVn stars have been developed by Smak (1983),

Cannizzo (1984), and Tsugawa, Osaki (1997). To explain the peculiar outburst behavior in EG Cnc, a modification of the disk-instability model for hydrogen-rich dwarf novae was introduced by Osaki et al. (2001), adding an idea of the viscosity decay in the cold disk. A similar idea may be necessary in reconstructing the full outburst property of 2003aw, although we need further observations to clarify the total outburst pattern including that outside the superoutbursts¹. The repetition of the large flares after the plateau phase will be a key also in trying to consider the behavior of 2003aw with the models making use of the variation in the mass-transfer rate (Warner 1995c), or the effect of the magnetic fields on the primary white dwarf (Woudt, Warner 2003).

The repetitive-outburst (high) state casts another problem concerning a notable difference in the superoutburst between 2003aw and EG Cnc. In 2003aw, small brightness variations with an amplitude of ≥ 0.4 mag and a period of ~ 16 h were observed, other than a large flare of 1.9 mag by Woudt, Warner (2003), while corresponding modulations were not reported in EG Cnc (Patterson et al. 1998; Kato et al. 2004a). The resemblance of the outburst-light curves between an AM CVn star and a WZ Sge star was first pointed out by Kato et al. (2004b) concerning V803 Cen and WZ Sge. After the initial, long outburst stage, V803 Cen showed ~ 1 mag oscillations with periods of 0.8–1.0 d, which resembles the rebrightening stage of the 2001 superoutburst in WZ Sge. Kato et al. (2004b) attributed those oscillations to some sort of thermal disk-instability, instead of the full-disk outbursts. To fully understand the outburst mechanism in 2003aw, we must explain the 0.4-mag oscillations with a timescale of 16 h and the occasional flares up to 2 mag.

The authors are thankful to amateur observers for continuously reporting their valuable observations to the VSNET. We are indebted to P. A. Woudt for his notification of the outburst and useful comments on the draft as the referee. This work is supported by Research Fellowships of Japan Society for the Promotion of Science for Young Scientists (MU and RI), and Grants-in-Aid for the 21st Century COE “Center for Diversity and Universality in Physics” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, and also by Grants-in-Aid from MEXT (No. 13640239, 15037205).

References

- Cannizzo, J. K. 1984, *Nature*, 311, 443
- Chornock, R., & Fillipenko, A. V. 2003, *IAU Circ.*, 8084
- Fernie, J. D. 1989, *PASP*, 101, 225
- Jha, S., Garnavich, P., Challis, P., Kirshner, R., & Berlind, P. 1998, *IAU Circ.*, 6983
- Kato, T., Nogami, D., Matsumoto, K., & Baba, H. 2004a, *PASJ*, 56, S109
- Kato, T. et al. 2001, *IBVS*, 5120

¹ Note that the outburst pattern in AM CVn stars can vary in a timescale of years (Patterson et al. 2000; Kato et al. 2001; Kato et al. 2004b).

- Kato, T., Stubbings, R., Monard, B., Butterworth, N. D., Bolt, G., & Richards, T. 2004b, PASJ, 56, S89
- Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2004c, PASJ, 56, S1
- Osaki, Y., Meyer, F., & Meyer-Hofmeister, E. 2001, A&A, 370, 488
- Patterson, J. et al. 1998, PASP, 110, 1290
- Patterson, J., Walker, S., Kemp, J., O'Donoghue, D., Bos, M., & Stubbings, R. 2000, PASP, 112, 625
- Roelofs, G. H. A., Groot, P. J., Steeghs, D., Nelemans, G. 2004, Rev. Mexicana Astron. Astrof., 20, 254
- Smak, J. 1983, Acta Astronomica, 33, 333
- Solheim, J.-E. 1995, Baltic Astronomy, 4, 363
- Stellingwerf, R. F. 1978, ApJ, 224, 953
- Tsugawa, M., & Osaki, Y. 1997, PASJ, 49, 75
- Warner, B. 1995a, Ap&SS, 225, 249
- Warner, B. 1995b, Cataclysmic Variable Stars (Cambridge: Cambridge University Press)
- Warner, B. 1995c, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle & M. Orio (Dordrecht: Kluwer Academic Publishers), 325
- Wood-Vasey, W. M., Aldering, G., Nugent, P., & Li, K. 2003, IAU Circ., 8077
- Woudt, P. A., & Warner, B. 2003, MNRAS, 345, 1266
- Zacharias, N., Urban, S. E., Zacharias, M. I., Wycoff, G. L., Hall, D. M., Monet, D. G., & Rafferty, T. J. 2004, AJ, 127, 3043